arXiv:1211.5427v1 [math.NA] 23 Nov 2012

Finite difference schemes on quasi-uniform grids for Bvps on infinite intervals

Riccardo Fazio and Alessandra Jannelli
Department of Mathematics, University of Messina
Viale F. Stagno D'Alcontres 31, 98166 Messina, Italy
rfazio@unime.it ajannelli@unime.it

December 5, 2011

Abstract

The classical numerical treatment of boundary value problems defined on infinite intervals is to replace the boundary conditions at infinity by suitable boundary conditions at a finite point, the so-called truncated boundary. A truncated boundary allowing for a satisfactory accuracy of the numerical solution has to be determined by trial and errors and this seems to be the weakest point of the classical approach. On the other hand, the free boundary approach overcomes the need for a priori definition of the truncated boundary. In fact, in a free boundary formulation the unknown free boundary can be identified with a truncated boundary and being unknown it has to be found as part of the solution.

In this paper we consider a different way to overcome the introduction of a truncated boundary, namely finite differences schemes defined on quasi-uniform grids. A quasi-uniform grid allows us to describe the infinite domain by a finite number of intervals. The last node of such grid is placed on infinity so right boundary conditions are taken into account exactly. We apply the proposed approach to the Falkner-Skan model and to a problem of interest in foundation engineering. The obtained numerical results are found in good agreement with those available in literature.

Key Words. non linear boundary value problems, infinite intervals, quasi-uniform grid, finite difference methods.
AMS Subject Classifications. 65L10, 65L12, 34B15.
Acknowledgement. The research of this work was partially supported by the University of Messina.

1 Introduction

The classical numerical treatment of boundary value problems (Bvps) on infinite intervals is to replace the original problem by one defined on a finite interval, where a finite value, the so-called truncated boundary, is used instead of infinity (see, for instance, Collatz [6, pp. 150-151] or Fox [14, p. 92]). For the accepted numerical solution, the value of the truncated boundary is varied until the computed results stabilize, at least, to a prefixed number of significant digits. However, a truncated boundary allowing for a satisfactory accuracy of the numerical solution has to be determined by trial and errors and this seems to be the weakest point of the classical approach. Hence, a priori definition of the truncated boundary was indicated by Lentini and Keller [17] as an important area of research.

A theory for defining asymptotic boundary conditions to be imposed at the truncated boundary has been developed by de Hoog and Weiss [7], Lentini and Keller [17] and Markowich [20, 21]. The asymptotic boundary conditions have been applied successfully to the numerical approximation of the so-called “connecting orbits” problems of dynamical systems, see Beyn [3, 4, 5]. Those problems are of interest, not only in connection with dynamical systems, but also in the study of traveling wave solutions of partial differential equations of parabolic and hyperbolic type as shown by Beyn [4], Friedman and Doedel [15], Bai et al. [2], and Liu et al. [19].

A free boundary formulation was proposed by Fazio [9] where the unknown free boundary was identified with a truncated boundary. In this approach the free boundary is unknown and has to be found as part of the solution. This free boundary approach overcomes the need for a priori definition of the truncated boundary. The free boundary formulation has been applied to: the Blasius problem [9], the

Falkner-Skan model [10], a model describing the flow of an incompressible fluid over a slender parabola of revolution [11], a connecting orbit problem [12], and a problem in foundation engineering [13].

Here we consider finite differences schemes on quasi-uniform grids applied to the numerical solution of BVPs defined on infinite intervals. This is a different way to avoid the definition of a truncated boundary. Quasi-uniform grids have successfully applied to the numerical solution of partial differential equations on unbounded domains, see Koleva [16] and the references quoted therein. In the following sections we consider two test problems. The first is the Falkner-Skan model of boundary layer theory. The last one is a problem of interest in foundation engineering.

We notice that the free boundary approach or a quasi-uniform grid strategy are as simple as the classical truncated boundary one in contrast with the asymptotic boundary approach, in this context see also the opinion expressed by J. R. Ockendon [23].

In the last section, we point out some conclusions supported by the evidences of the present work and indicate a possible way to extend the proposed approach to Bvps defined on the whole real line.

2 Finite differences on quasi-uniform grids

Let us consider the smooth strict monotone quasi-uniform map $x = x(\xi)$, the so-called grid generating function,

$$x = -c \cdot \ln(1 - \xi) \ , \tag{2.1}$$

where $\xi \in [0, 1]$, $x \in [0, \infty]$, and $c > 0$ is a control parameter. We notice that $x_{N-1} = c \ln N$ for (2.1). The problem under consideration can be discretized by introducing a uniform grid ξ_n of $N+1$ nodes in $[0, 1]$ with $\xi_0 = 0$ and $\xi_{n+1} = \xi_n + h$ with $h = 1/N$, so that x_n is a quasi-uniform grid in $[0, \infty]$. The last interval in (2.1), namely $[x_{N-1}, x_N]$, is infinite but the point $x_{N-1/2}$ is finite, because the non integer nodes are defined by

$$x_{n+\alpha} = x\left(\xi = \frac{n+\alpha}{N}\right) \ ,$$

with $n \in \{0, 1, \ldots, N-1\}$ and $0 < \alpha < 1$. This map allows us to describe the infinite domain by a finite number of intervals. The last node of such grid is placed on infinity so right boundary condition is taken into account correctly.

Figure 1 shows the quasi-uniform mesh $x = x_n$, $n = 0, 1, \ldots, N$ defined by (2.1) with $c = 5$ and $N = 20$.



Figure 1: *A quasi-uniform mesh* $x = x_n$. *We notice that the last mesh-point is* $x_N = \infty$.

We can define the values of $u(x)$ on the mid-points of the grid

$$u_{n+1/2} \approx \frac{x_{n+1} - x_{n+1/2}}{x_{n+1} - x_n} u_n + \frac{x_{n+1/2} - x_n}{x_{n+1} - x_n} u_{n+1} \ . \tag{2.2}$$

As far as the first derivative is concerned we can apply the following approximation

$$\left. \frac{du}{dx} \right|_{n+1/2} \approx \frac{u_{n+1} - u_n}{2\left(x_{n+3/4} - x_{n+1/4}\right)} \ . \tag{2.3}$$

This formulae uses the value $u_N = u_\infty$, but not $x_N = \infty$. In order to justify finite difference formula (2.3) we note that, by considering $u = u(\xi(x))$, we can write

$$\left. \frac{du}{dx} \right|_{n+1/2} = \left. \frac{du}{d\xi} \right|_{n+1/2} \left. \frac{d\xi}{dx} \right|_{n+1/2} \approx \frac{u_{n+1} - u_n}{\xi_{n+1} - \xi_n} \frac{2\left(\xi_{n+3/4} - \xi_{n+1/4}\right)}{2\left(x_{n+3/4} - x_{n+1/4}\right)} \ . \tag{2.4}$$

The last formula on the right hand side of equation (2.4) reduces to the right hand side of equation (2.3) because we are using a uniform grid for ξ and therefore $2\left(\xi_{n+3/4} - \xi_{n+1/4}\right) = \xi_{n+1} - \xi_n$.

The two finite difference approximations (2.2) and (2.3) have order of accuracy $O(N^{-2})$.

3 Bvps on infinite intervals

Let us consider the class of Bvps defined on an infinite interval

$$\begin{aligned} & \frac{d\mathbf{u}}{dx} = \mathbf{f}\left(x,\mathbf{u}\right) \ , \quad x \in [0,\infty) \ , \\ & \mathbf{g}\left(\mathbf{u}(0),\mathbf{u}(\infty)\right) = \mathbf{0} \ , \end{aligned} \tag{3.1}$$

where $\mathbf{u}(x)$ is an $d-$dimensional vector with $u_\ell(x)$ for $\ell = 1,\ldots,d$ as components, $\mathbf{f} : [0,\infty) \times \mathbb{R}^d \rightarrow \ \mathbb{R}^d$, and $\mathbf{g} : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.

A finite difference scheme on a quasi-uniform grid for the class of Bvps (3.1) can be defined by using the approximations given by (2.2) and (2.3). We denote by the $d-$dimensional vector $\mathbf{U}_n$ the numerical approximation to the solution $\mathbf{u}(x_n)$ of (3.1) at the points of the mesh, that is for $n = 0,1,\ldots,N$. A second order finite difference scheme for (3.1) can be written as follows:

$$\begin{gathered} \mathbf{U}_{n+1} - \mathbf{U}_n - a_{n+1/2}\mathbf{f}\left(x_{n+1/2}, b_{n+1/2}\mathbf{U}_{n+1} + c_{n+1/2}\mathbf{U}_n\right) = \mathbf{0} \ , \\ \text{for} \quad n = 0,1,\ldots,N-1 \\ \mathbf{g}\left(\mathbf{U}_0,\mathbf{U}_N\right) = \mathbf{0} \ , \end{gathered} \tag{3.2}$$

where

$$\begin{aligned} a_{n+1/2} &= 2\left(x_{n+3/4} - x_{n+1/4}\right) \ , \\ b_{n+1/2} &= \frac{x_{n+1/2} - x_n}{x_{n+1} - x_n} \ , \\ c_{n+1/2} &= \frac{x_{n+1} - x_{n+1/2}}{x_{n+1} - x_n} \ , \end{aligned} \tag{3.3}$$

for $n = 0,1,\ldots,N-1$.

It is evident that (3.2) is a nonlinear system of $d \cdot (N+1)$ equations in the $d \cdot (N+1)$ unknowns $\mathbf{U} = (\mathbf{U}_0,\mathbf{U}_1,\ldots,\mathbf{U}_N)^T$. We notice that $b_{n+1/2} \approx c_{n+1/2} \approx 1/2$ for all $n = 0,1,\ldots,N-2$, but when $n = N-1$, then $b_{N-1/2} = 0$ and $c_{N-1/2} = 1$. On the contrary, we choose to set $b_{N-1/2} = b_{N-3/2}$ and $c_{N-1/2} = c_{N-3/2}$ in order to avoid a suddenly jump for the coefficients of (3.2). As it will be clear from the results reported in the next section this produces a much smaller error in the numerical solution of the system at x_N.

For the solution of (3.2) we can apply the classical Newton's method along with the simple termination criterion

$$\frac{1}{d(N+1)}\sum_{\ell=1}^{d}\sum_{n=0}^{N}|\Delta U_{n\ell}| \leq \mathrm{TOL}\ ,$$

where $\Delta U_{n\ell}$, for $n = 0,1,\ldots,N$ and $\ell = 1,2,\ldots,d$, is the difference between two successive iterate components and TOL is a fixed tolerance. The results listed in the next sections were computed by setting $\mathrm{TOL} = 1\mathrm{E}-6$.

4 The Falkner-Skan model

The Falkner-Skan model [8], of boundary layer theory [25], is given by

$$\begin{aligned}
&\frac{d^3u}{dx^3} + u\frac{d^2u}{dx^2} + P\left[1-\left(\frac{du}{dx}\right)^2\right] = 0 \\
&u(0) = \frac{du}{dx}(0) = 0, \qquad \frac{du}{dx}(\infty) = 1\ ,
\end{aligned} \tag{4.1}$$

is a Bvp defined on a semi-infinite interval.

As a first step we rewrite the Falkner-Skan equation in (4.1) as a first order system by setting

$$u_{i+1}(x) = \frac{d^iu}{dx^i}(x)\ , \quad \text{for } i = 0,1,2\ .$$

In this way the original Bvp (4.1) specializes to

$$\begin{aligned}
&\frac{du_1}{dx} = u_2 \\
&\frac{du_2}{dx} = u_3 \\
&\frac{du_3}{dx} = -u_1u_3 - P(1-{u_2}^2) \\
&u_1(0) = u_2(0) = 0\ , \qquad u_2(\infty) = 1\ ,
\end{aligned} \tag{4.2}$$

that is,

$$\begin{aligned}
\mathbf{u} &= (u_1,u_2,u_3)^T \\
\mathbf{f}(x,\mathbf{u}) &= \left(u_2,u_3,u_4,-u_1u_3-P(1-{u_2}^2)\right)^T \\
\mathbf{G}\left(\mathbf{u}(0),\mathbf{u}(\infty)\right) &= \left(u_1(0),u_2(0),u_2(\infty)-1\right)^T
\end{aligned}$$

in (3.1).

4.1 Numerical results

Some of the numerical results obtained for the test problem introduced in this section are reported in this subsection. Figure 2 shows the numerical solution of Falkner-Skan model (4.1) with $P = 1$.



Figure 2: *Numerical solution of Falkner-Skan model with* $P = 1$ *obtained with the map* $x = x(\xi)$ *defined by (2.1) with* $c = 5$ *for* $N = 80$.

Table 1 lists the numerical approximations of $\frac{d^2u}{dx^2}(0)$ and $\frac{d^2u}{dx^2}(\infty)$ for increasing values of N. For all values of N we used the initial iterate

$$u_1(x) = u_2(x) = 1/2 \ , \quad u_3(x) = 10^{-2} \ .$$

Let us remark here that the same values of $\frac{d^2u}{dx^2}(0)$ were obtained by using $b_{N-1/2} = 0$ and $c_{N-1/2} = 1$ in (3.2), but, on the contrary, larger values of $\frac{d^2u}{dx^2}(\infty)$ were computed. In table 2 we compare the obtained numerical results with those available in literature: the agreement is really good.

Table 1: *Numerical approximation of* $\frac{d^2u}{dx^2}(0)$ *and* $\frac{d^2u}{dx^2}(\infty)$.

N	iter	$\frac{d^2u}{dx^2}(0)$	$\frac{d^2u}{dx^2}(\infty)$
20	6	1.238724	$-0.21 \cdot 10^{-7}$
40	5	1.234124	$0.24 \cdot 10^{-7}$
80	5	1.232972	$-0.33 \cdot 10^{-7}$
160	5	1.232684	$0.14 \cdot 10^{-7}$
320	5	1.232612	$-0.25 \cdot 10^{-7}$
640	5	1.232594	$0.39 \cdot 10^{-7}$
1280	5	1.232589	$0.33 \cdot 10^{-7}$

Table 2: *Comparison of* $\frac{d^2u}{dx^2}(0)$ *and free or truncated boundary (*x_ε *and* x_∞ *respectively) for the Homann (*$P = 1/2$*) and Hiemenz (*$P = 1$*) flows.*

	This paper		Fazio [10]		Nasr et al. [22]		Asaithambi [1]	
	Quasi-uniform		Free BF		Chebyshev method		Finite difference	
P	x_N	$\frac{d^2u}{dx^2}(0)$	x_ε	$\frac{d^2u}{dx^2}(0)$	x_∞	$\frac{d^2u}{dx^2}(0)$	x_∞	$\frac{d^2u}{dx^2}(0)$
0.5					3.7	0.927805		
0.5	∞	0.927681	5.09	0.927680	7.4	0.927680	5.67	0.927682
1					3.5	1.232617		
1	∞	1.232589	5.19	1.232588	7.	1.232588	5.14	1.232589

5 A pile in soil

Here we consider a problem that was already used by Lentini and Keller [17] to test the asymptotic boundary conditions approach. This problem is of special interest here because none of the solution components is a monotone function, on $[0,\infty)$, see [12, 13]. Let $u(x)$ be the deflection of a semi-infinite pile embedded in soft soil at a distance x below the surface of the soil. The governing differential equation for the movement of the pile, in dimensionless form, is given by:

$$\begin{gathered}
\frac{d^4u}{dx^4} = -P_1\left(1-e^{-P_2u}\right) \ , \qquad x \in [0,\infty) \ , \\
\frac{d^2u}{dx^2}(0) = 0 \ , \qquad \frac{d^3u}{dx^3}(0) = P_3 \ , \qquad u(\infty) = \frac{du}{dx}(\infty) = 0 \ ,
\end{gathered} \tag{5.1}$$

where P_1 and P_2 are positive material constants. As far as the boundary conditions are concerned, at the origin a zero moment and a positive shear P_3 are assumed and from physical considerations it follows that $u(x)$ and all its derivatives go to zero at infinity, so that, the zero asymptotic boundary conditions can be imposed. This problem is of interest in foundation engineering: for instance, in the design of drilling rigs above the ocean floor.

The governing differential equation in (5.1) can be rewritten as a first order system by setting:

$$u_{i+1}(x) = \frac{d^iu}{dx^i}(x) \ , \quad \text{for } i = 0,1,2,3 \ .$$

In this way the original Bvp (5.1) specializes to

$$\begin{aligned}
&\frac{du_1}{dx} = u_2 \ , \\
&\frac{du_2}{dx} = u_3 \ , \\
&\frac{du_3}{dx} = u_4 \ , \\
&\frac{du_4}{dx} = -P_1\left(1-e^{-P_2u_1}\right) \ , \\
&u_3(0) = 0 \ , \qquad u_4(0) = P_3 \ , \qquad u_1(\infty) = 0 \ , \qquad u_2(\infty) = 0 \ ,
\end{aligned} \tag{5.2}$$

that is,

$$\mathbf{u} = (u_1, u_2, u_3, u_4)^T$$

$$\mathbf{f}(x,\mathbf{u}) = \left(u_2, u_3, u_4, -P_1\left(1-e^{-P_2 u_1}\right)\right)^T$$
$$\mathbf{G}\left(\mathbf{u}(0),\mathbf{u}(\infty)\right) = \left(u_3(0), u_4(0)-P_3, u_1(\infty), u_2(\infty)\right)^T$$

in (3.1).

5.1 Numerical results

In this subsection we report some of the numerical results obtained for the test problem introduced in this section. In order to be able to compare our numerical results we used the same parameter values employed by Lentini and Keller [17]

$$P_1 = 1, \quad P_2 = \frac{1}{2} \quad \text{and} \quad P_3 = \frac{1}{2} \; .$$

Moreover, we choose to consider the values of the missing initial conditions $u_1(0)$ and $u_2(0)$ as representative results.

Figure 3 shows the numerical solution of the Bvp (5.1).

Table 3 lists the numerical approximations of $u(0)$ and $\frac{du}{dx}(0)$ for increasing values of N. For all values of N we used the initial iterate

$$u_1(x) = u_2(x) = u_3(x) = u_4(x) = 1 \; .$$

Table 3: *Numerical approximation of* $u(0)$ *and* $\frac{du}{dx}(0)$.

N	iter	$u(0)$	$-\frac{du}{dx}(0)$
20	5	1.420337	0.807289
40	5	1.421243	0.807934
80	5	1.421469	0.808094
160	5	1.421526	0.808135
320	5	1.421540	0.808145
640	5	1.421544	0.80815
1280	5	1.421544	0.80815



Figure 3: *Numerical solution of pile model (5.1) obtained with the map* $x = x(\xi)$ *defined by (2.1) with* $c = 5$ *for* $N = 80$.

Table 4: *Comparison of* $u(0)$, $\dfrac{du}{dx}(0)$ *and free or truncated boundary (*x_ε *and* x_∞ *respectively) for the pile problem.*

This paper		Fazio [10]		Lentini and Keller [17]	
Quasi-uniform $x_N = \infty$		Free BF $x_\varepsilon = 17.75$		Asymptotic BCs $x_\infty = 10$	
$u(0)$	$\dfrac{du}{dx}(0)$	$u(0)$	$\dfrac{du}{dx}(0)$	$u(0)$	$\dfrac{du}{dx}(0)$
1.421544	-0.808145	1.42154	-0.808144	1.4215	-0.80814

As far as the missing initial conditions for the pile problem are concerned, a comparison of the obtained values has been considered in table 4. It is easily seen that our results are in good agreement with those available in literature. Lentini and Keller [17] used the mentioned asymptotic boundary conditions and employed PASVAR, a routine based upon the trapezoidal difference scheme with automatic mesh refinement and deferred corrections as described by Lentini and Pereyra [18]. That software is rather sophisticated because it adjusts automatically the mesh and the order of accuracy of the method employed. Fazio [10] used a free boundary formulation of the pile problem as mentioned in the introduction and the Keller's box finite difference method.

6 Concluding remarks

The numerical results for the test problems reported in the two previous sections show that finite difference schemes on quasi-uniform grids are an effective way to solve Bvps defined on infinite intervals. The application of finite difference schemes on quasi-uniform grids overcomes the need for a priori definition of the truncated boundary.

A further advantage in using a family of quasi-uniform grids in calculations is the possibility to improve the obtained accuracy. The algorithm is the same for many finite difference methods: for numerical differentiation or integration, solving systems of ordinary or partial differential equations. It is based on Richardson extrapolation, introduced by Richardson in [24]. We carry out two calculations on embedded uniform or quasi-uniform grids with total number of nodes N and $2N$. All nodes of largest steps are identical to even nodes of denser grid due to uniformity. Suppose that we use numerical method with order of accuracy p in order to find an approximation of a scalar value T. For smooth enough solutions the error can be decomposed into a sum of inverse powers of N. The Richardson formula

$$R_{2N} = \frac{T_{2N} - T_N}{2^p - 1} \tag{6.1}$$

defines the main term of such sum. This formula is asymptotically exact when $N \to \infty$ if we use uniform or quasi-uniform grids. Hence, it gives the real value of numerical solution error without knowledge of exact solution. We can apply (6.1) as a single-step correction formula

$$T = T_{2N} + R_{2N} + O(N^{-p-s}) \tag{6.2}$$

and increase the order of accuracy of our approximation. In general, we have $s = 2$ in (6.2) for a symmetrical finite difference scheme, but only $s = 1$ for non-symmetrical ones. Such enlargement of accuracy requires to solve the same problem twice and only few further arithmetical operations and so it is very cheap.

I could be possible to apply Richardson extrapolation to the results reported in table 1 and in table 3. In this way we can avoid to solve the given problem with a large number of grid-points.

Let us discuss, at the end of this work, a possible way to extend the finite difference schemes on quasi-uniform grids to the numerical solutions of problems defined on the whole real line, for instance, the connecting orbits problems mentioned in the introduction. For these problems, all boundary conditions are imposed at plus or minus infinity. In such a case it is possible to use the tangential quasi-uniform grid

$$x_n = c \cdot \tan\left(\frac{n\pi}{2N}\right) ,$$

where $c > 0$ is a control parameter. In fact, if $n = -N, -N+1, \ldots -1, 0, 1, \ldots, N-1, N$, then this tangential grid cover the whole infinite line, and in particular we have that $x_{-N} = -\infty$.

References

[1] A. Asaithambi, A finite-difference method for the solution of the Falkner-Skan equation, Appl. Math. Comput. 92 (1998) 135–141.

[2] F. Bai, A. Spence, A. M. Stuart, The numerical computation of heteroclinic connections in systems of gradient partial differential equations, SIAM J. Appl. Math. 53 (1993) 743–769.

[3] W. J. Beyn, Global bifurcation and their numerical computation, in: D. Rossed, B. D. Dier, A. Spence (eds.), Bifurcation: Numerical Techniques and Applications, Kluwer, Dordrecht, 1990, pp. 169–181.

[4] W. J. Beyn, The numerical computation of connecting orbits in dynamical systems, IMA J. Numer. Anal. 9 (1990) 379–405.

[5] W. J. Beyn, Numerical methods for dynamical systems, in: W. Light (ed.), Advances in Numerical Analysis, Clarendon Press, Oxford, 1992, pp. 175–236.

[6] L. Collatz, The Numerical Treatment of Differential Equations, 3rd ed., Springer, Berlin, 1960.

[7] F. R. de Hoog, R. Weiss, An approximation theory for boundary value problems on infinite intervals, Computing 24 (1980) 227–239.

[8] V. M. Falkner, S. W. Skan, Some approximate solutions of the boundary layer equations, Philos. Mag. 12 (1931) 865–896.

[9] R. Fazio, The Blasius problem formulated as a free boundary value problem, Acta Mech. 95 (1992) 1–7.

[10] R. Fazio, The Falkner-Skan equation: numerical solutions within group invariance theory, Calcolo 31 (1994) 115–124.

[11] R. Fazio, A novel approach to the numerical solution of boundary value problems on infinite intervals, SIAM J. Numer. Anal. 33 (1996) 1473–1483. URL `http://epubs.siam.org/sam-bin/dbq/toc/SINUM/33/4`

[12] R. Fazio, A survey on free boundary identification of the truncated boundary in numerical BVPs on infinite intervals, J. Comput. Appl. Math. 140 (2002) 331–344.

[13] R. Fazio, A free boundary approach and Keller's box scheme for BVPs on infinite intervals, Int. J. Computer Math. 80 (2003) 1549–1560.

[14] L. Fox, Numerical Solution of Two-point Boundary Value Problems in Ordinary Differential Equations, Clarendon Press, Oxford, 1957.

[15] M. J. Friedman, E. J. Doedel, Numerical computation and continuation of invariant manifolds connecting fixed points, SIAM J. Numer. Anal. 28 (1991) 789–808.

[16] M. N. Koleva, Numerical solution of the heat equation in unbounded domains using quasi-uniform grids, in: I. Lirkov, S. Margenov, J. Wasniewski (eds.), Large-scale Scientific Computing, vol. 3743 of Lecture Notes in Comput. Sci., 2006.

[17] M. Lentini, H. B. Keller, Boundary value problems on semi-infinite intervals and their numerical solutions, SIAM J. Numer. Anal. 17 (1980) 577–604.

[18] M. Lentini, V. Pereyra, An adaptive finite difference solver for two-point boundary value problems with mild boundary layers, SIAM J. Numer. Anal. 14 (1977) 91–111.

[19] L. Liu, G. Moore, R. D. Russell, Computation and continuation of homoclinic and heteroclinic orbits with arclength parameterization, SIAM J. Sci. Comput. 18 (1997) 69–93.

[20] P. A. Markowich, A theory for the approximation of solution of boundary value problems on infinite intervals, SIAM J. Math. Anal. 13 (1982) 484–513.

[21] P. A. Markowich, Analysis of boundary value problems on infinite intervals, SIAM J. Math. Anal. 14 (1983) 11–37.

[22] H. Nasr, I. A. Hassanien, H. M. El-Hawary, Chebyshev solution of laminar boundary layer flow, Int. J. Comput. Math. 33 (1990) 127–132.

[23] J. R. Ockendon, Math. Rev., 84c:34021.

[24] L. F. Richardson, The deferred approach to the limit, Phil.Trans., A 226 (1927) 299–349.

[25] H. Schlichting, K. Gersten, Boundary Layer Theory, 8th ed., Springer, Berlin, 2000.